

100-11178

Library



West Virginia University

EVANS DALE LIBRARY
WEST VIRGINIA
UNIVERSITY

West Virginia University Libraries



3 0802 100341440 8

West Virginia University Library

This book is due on the date indicated below.

APR 24 1966

APR 4 1966

SEP 20 1973

MAY 20 1961

2 JUN 82 2 0 DEC 82

RETURNED



VOLUME IV.


NUMBER 4.

BULLETIN 39.

West Virginia Agricultural Experiment Station.

MORGANTOWN, W. VA.

VEGETABLES.



Buckeye State.

By F. WM. RANE.

JANUARY, 1895.



CHARLESTON, W. VA.
MOSES W. DONNALLY, PUBLIC PRINTER
1895.

Board of Regents of the West Virginia University.

District.	Name of Regent.	P. O. Address.
1.	J. B. SOMMERVILLE,	Wheeling.
2.	CLARENCE L. SMITH,	Fairmont.
3.	J. S. WITHERS,	Glenville.
4.	JOHN BASSEL,	Clarksburg.
5.	J. M. HOLSWADE,	Spencer.
6.	JAMES H. STEWART,	Raymond City.
7.	WIRT A. FRENCH,	Princeton.
8.	M. J. KESTER,	Union.
9.	J. F. BROWN,	Charleston.
10.	THOMAS J. FARNSWORTH,	Buckhannon.
11.	JOSEPH MORELAND,	Morgantown.
12.	JOHN A. ROBINSON,	Patterson's Depot.
13.	DR. W. W. BROWN,	Kabletown.

Members of the Station Committee.

JOHN A. ROBINSON.	C. L. SMITH,
J. S. WITHERS,	JAS. H. STEWART.
DR. W. W. BROWN.	

<i>Prest. of Board of Regents,</i>	-	-	JOHN A. ROBINSON.
<i>Acting President of the University,</i>	-	-	P. B. REYNOLDS, LL. D.
<i>Treasurer,</i>	-	-	A. R. WHITEHILL, PH. D.

Station Staff.

JOHN A. MYERS, PH. D.,	-	Director.
F. WM. RANE, M. S.	-	Horticulturist and Microscopist.
A. D. HOPKINS, PH. D.,	-	Entomologist.
RUDOLF DE ROODE, PH. D.,	-	Chemist.
W. E. RUMSEY, B. S. AGR.,	-	Assistant Entomologist.
M. ELIZABETH HEERMANS,	-	Stenographer.

HORTICULTURAL DEPARTMENT.

Bulletin No. 39. Vegetables.

CONTENTS.

	Pages
Introductory	138
Trial Grounds	138
Beans, Bush	(Illustrated) 139
Beans, Bush Lima	140
Beans, Pole	140
Beans, Pole Lima	140
Beets	141
Cauliflower	142
Cabbage, First Early	(Illustrated) 142
Cabbage, Second Early	144
Cabbage, Late	144
Celery	(Illustrated) 144
Cucumbers	146
Egg Plant	146
Kale	(Illustrated) 146
Kohl-Rabi	147
Lettuce	147
Miscellaneous Végétables	155
Musk Melon	148
Onions	(Illustrated) 148
Peas	149
Pepper	150
Radish	150
Salsify	151
Squash	151
Sweet Corn	151
Tomato	152
Water Melon	155
Donations to Horticultural Department	155
Remarks	156
Macrosporium of the Pepper	150
Flea Beetle (Black) on Tomato	152

VARIETY TESTS OF VEGETABLES.

By F. WM. RANE, M. S.

Olericulture, or vegetable gardening, is progressive, and to keep abreast of the times some little attention must be paid to it. During the season of 1894, the Horticultural Department entered upon this line of work for the first time.

The testing of the seeds of new varieties which seedsmen and private individuals are continually offering in their attempts to find something better than we already have, is an important work, and one that requires labor and expense as well. In itself it is a very simple matter, but the average farmer and gardener cannot afford to spend his time and money on an extensive test.

To aid the gardener, then, the Station undertakes this work. Heretofore our experiments have been conducted on a limited scale, owing to lack of available ground. We have tested those seeds which we have been requested to test and have reported to the sender, but have not published the results.

Our purpose is to eliminate the undesirable new varieties, and to establish the worth of the promising ones. This being our first year, we selected many of the older and more familiar varieties than we shall hereafter, in order that every one might be able to compare those grown by themselves with the new introductions. The coming year we will retain only those showing superior points for further trial, to which we will add this year's introductions.

The Trial Grounds.

January, 1894, the Director turned the land formerly used by the late Agriculturist, over to this Department. It contains about three acres, is located just across the street from the Station building, and extends east and west, sloping from north-east to south-west. The soil is a reddish clay loam and has been used for raising corn, sorghum and tobacco, for the past few years.

After it was placed in our charge it was plowed, limed heavily, manured, and again plowed, harrowed, and gone over with the garden rake, to put it in order. Although a great amount of labor was spent on the land, it is needless to say that it did not take the place of a fertile garden, and all who read this bulletin must, therefore, take this fact into account. Not only was the soil new for gardening purposes, but we also had an extremely dry season. We know that our yield is below the average in quantity, but it is sufficient to establish the comparative merits or demerits of the various varieties of vegetables, which was our main object.

The following data were mainly taken under personal direction by Messrs. J. M. Johnson and A. J. Ireland, two promising young men in the Horticultural Department of our University.



Cut 2.—Bush Beans.

Bush Beans.

There are numerous varieties of bush beans, and the variety to be selected depends upon the use you desire to make of them. They vary as to earliness, productiveness, color, size and quality. The bean can be grown on almost any soil but revels best in those comparatively light and well-drained. Never plant too early, as they are liable to be injured by the cool damp weather in early spring. The time recommended as safest is at corn planting, and by planting every two or three weeks from this time until the end of July, a succession may be had throughout the year. The varieties named in the following table were planted in rows, three and one-half feet apart, each variety occupying seventeen feet in the row. The seed was sown on May 8th. The pods were picked and weighed as they ripened; also other data taken. In cut 2 are found representative specimens.

Number.....	VARIETY.	Seedsman.	Date of Edible Ma- turity.	No. Plants matured	Height.	Diameter of Head.	Total Yield.	Av. number beans per pod.
1	China Red Eye Bush.....	Landreth..	July 6...	56	13	13	5.7	5
2	E'y Yellow Six Week.....	Landreth..	6...	66	15	14	11.5	5
3	E'y Brown Six Week.....	Landreth..	6...	63	13	12	8.3	5
4	Ex. Ey Red Valentine.....	Landreth..	6...	63	14	12	9.9	5
5	Landreth's "First in Market".	Landreth..	31...	30	14	11	3.1	7
6	Dwf. German Wax.....	Landreth..	6...	68	14	16	9.	6
7	Dwf. Kidney Wax.....	Landreth..	7...	19	13	11	3.5	5
8	White Wax Bush.....	Landreth..	13...	30	11	11	4.2	4
9	Landreth's Scarlet Bean.....	Landreth..	17...	19	13	13	3.5	6
10	Pink Eye Wax.....	Landreth..	10...	30	12	10	5.2	6
11	Golden Eyed Wax.....	Landreth..	13...	45	15	14	6.7	5
12	White Valentine Bush.....	Landreth..	7...	33	13	13	8.3	4½
13	White Cranberry Bush.....	Landreth..	31...	36	16	20	4.7	5
31	New Stringless Green Pod.....	Burpee....	7...	51	16	15	9.3	5
32	Speckled Wax.....	Barteldes..	31...	54	14	12	8.6	5
33	Eyed Golden Wax.....	Barteldes..	10...	47	16	14	9.1	4½
38	Bush Bean Round Yellow.....	Dreer.....	16...	52	13	14	8.3	5
39	Bush Bean Sion House.....	Dreer.....	Aug. 1...	54	14	23	7.5	5½
40	Thornburn's Ex. Ey. Refugee..	Dreer.....	July 23...	51	12	14	7.9	6
41	Thornburn's Ex. Ey. Market.....	Thornburn	Aug. 1...	19	12	17	8.4	4
42	Vick's Prolific Pickler.....	Vick Sons..	1...	41	21	24	8.6	4
43	Wisconsin Tree.....	J.A. Everitt	July 16...	47	14	20	9.8	4
44	Jackson Wonder.....	J.A. Everitt	Aug. 10...	39	12	14	12.3	7

Lima Beans.

The *Dwarf Limas* did not ripen well, and the percentage of germination was poor with some varieties. We give the following table, (see below) however, that our gardeners may compare them with their own.

Number.	VARIETY.	Seedsman.	Date of Ed- ible Maturity.	No. Plants Matured.	Height inches.	Diameter of Head inches.	Total Yie d Pounds.	Average No. beans per pod	Av. Weight per Plant, Ounces.	Remarks.
23	Burpee's Bush Lima.....	U. S. Dept....	Aug. 11..	34	13	12	7.3	5	3.36	About 1 quart. Not matured.
26	Dreer's Bush Lima.....	Johnson & Stokes....	Aug. 16..	6	10	18	4.5	4	12.00	Lay on ground. 3 quarts not matured.
27	Jackson's Wonder Dur- fee Lima	Johnson & Stokes....	Aug. 11....	29	13	18	11.6	5	6.40	Lay on ground.
28	Bush Lima.....	Burpee.....	Aug. 11....	13	10	18	4.7	5	4.56	Lay on ground.
34	Henderson's Dwarf Li- ma	N. B. G. Co..	Aug. 11....	36	14	13	8.4	5	3.68	Begn'g to run.
35	Kumerle Dwarf Lima....	N. B. G. Co..	Oct. 8.....	2	10	15	.9	4	7.20	Germinated poorly.
36	Buckbee's Dwarf Lima.	N. B. G. Co..	Aug. 11....	11	14	13	3.5	4	4.96	Three quarts. Not matured October 9.
37	Dreer's Dwarf Lima.....	N. B. G. Co..	Aug. 25....	10	13	12	3.4	5	5.44

The *Pole Limas* run more evenly. The column marked "Average per Hill" will give a fair, comparative yield for our soil and conditions.

Number.	VARIETY.	Seedsman.	Date of Ed- ible Maturity.	No. Hills Ma- ture .	Total Yield in Pounds.	Average per Hill.	Average No. Beans per Pod.
19	White Dutch or Case Knife.....	Landreth.	Aug. 2...	10	10	Lbs. Ozs. 1 0	5
29	Ky. Black Lima Pole Bean.....	Burpee....	Aug. 16	21	27.4	1 5 1/2	6
30	Horticultural Lima.....	Burpee.....	July 23...	10	17.3	1 11	5
44	Willow Leaf Lima.....	J.A. Everitt	Aug. 13...	11	12.7	1 13-5	4

Pole Beans.

The Pole Beans must be given more room than the dwarf. They are also tenderer and usually planted a fortnight later. Set the poles, when planting, so as not to disturb the young plants when they germinate. The varieties in the test were all planted May 8th. There were more hills of some varieties than of others, but in the column marked "Average per Hill," the comparisons can be easily seen.

Number.	VARIETY.	SEEDSMAN.	Date of Edible Matu- rity.	Number hills Matured.	Total Yield In Pounds.	Average Per Hill		Average No Beets Per Pod.
						lbs.	oz.	
15	Tall German Wax...	Landreth.....	Sept. 16	8	7.5		23-10	3
16	Caroline	Landreth.....	August 11	10	10.8	1	13-5	4
17	Giant Wax	Landreth.....	" 2	7	12.5	1	13	5
20	Southern Prolific ...	Landreth.....	" 3	16	7.9		8	7
21	Horticultural.....	Landreth.....	" 2	11	4.4		7	5
22	Fat Horse.....	Landreth.....	July 16	20	21.8	1	1½	6
24	Kentucky Wonder...	U. S. Dept.....	" 16	35	46.1	1	5½	8

Beets.

The varieties under test are mainly of the turnip-rooted forms, for the reason that these are in the greatest demand in the markets. There are many very valuable varieties at present. The best one to be grown will depend upon the fancy of the market as to color, size, shape, etc. The ground cannot be too rich nor too thoroughly pulverized. The roots generally bring the best prices early in the season, and it often happens that it pays better to harvest the crop when half grown, than to wait until fully matured. Like many other vegetables, they may be had continually by sowing in succession from two to three weeks apart. Always press the soil closely to the seed when planted, or in a dry time the seed will not germinate. The ground occupied by the early crop is always used for a second one after they are marketed. The majority of gardeners sow more seed than is necessary, and then thin out, leaving the desired number of plants to mature, using those thinned out as a substitute for spinach.

Number.	VARIETY.	Seedsman.	Total Yield	Weight. lbs.	Av. Wt., oz.	Av. Diam. in.	Character of Form.	Color of Sktn.
1	Dewey's Red Turnip	Landreth.....	20	2.5	1.60	1½	Long.	Dk. Red.
2	Eclipse.....	"	28	10.3	4.56	3	Conical.	Red.
3	Egyptian Ex. Ey. Turnip.....	"	17	1.5	1.60	2	Turnip.	Dk. Red.
4	Bassano Ex. Ey. Turnip.....	"	27	19	1.12	4	"	Bt. Red.
5	Ely Blood Red Turnip.....	"	15	3	23.20	2	Conical.	Ok. Red.
6	Imp. Blood Red Turnip.....	"	20	9.	7.20	3½	Turnip.	" "
7	Philadelphia Ey. Turnip.....	"	27	9.	4.96	3	"	" "
16	New Danish Imp. Sugar.....	Burpee.....	10	3.	4.80	2	Long.	Pale Red.
17	Columbia Be t.....	"	39	11.7	4.80	2½	Conical.	Pale Red.
18	Derigo Blood Turnip	U. S. Dept.....	23	18.	1.24	3	"	Red.
19	Burpee's Imp. Blood Turnip.....	Burpee	23	12	5.48	3	Turnip	Bt. Red.
20	Imp. Long Dk. Blood.....	Ferry.....	18	1.8	1.60	1	Long	Dk. Rwd.
21	Lentz.....	Barteldes.....	33	10.1	5.2	3	Conical.	Bt. Red.
22	Crosby's Egyptian.....	Dreer.....	35	11	8.28	2½	Turnip.	Red.
23	Crimson Chief Turnip.....	Johnson & Stokes;	28	13.8	8.00	3	"	Bt. Red.
24	Landreth's Very Early.....	Landreth.....	19	3.3	2.56	2	Conical.	" "
25	Lentz.....	"	23	9.5	6.40	2½	"	Red.
26	Bloodsdales H'f Round.....	"	34	14.	6.40	3	Hf. round	"
27	Bastian.....	"	17	5.5	4.80	2½	Conical.	Bt. Red.

Cauliflower.

Twelve varieties were selected for this test. Cauliflower demands a rich soil, in order to produce large heads. The plants were started in the greenhouse and after hardening off in the cold frame were transplanted out of doors. Twenty-five plants of each variety were grown. The best varieties as regards earliness and productiveness were the *Early Dwarf Erfurt*, H. A. March, and the *Alabaster*, Johnson & Stokes. These two ripened at the same time and were equally the largest heads. *Early Snowball*, Dreer, and *Model*, N. B. G. & Co., together with *Early Perfection*, March, were second best. The same cultivation that is given to cabbage will answer for cauliflower.



Cut 3.—First Early Cabbage.

First Early Cabbage.

Twenty-two varieties were used in the test of early cabbage. The plants, numbering twenty-five each variety, were transplanted into the garden April 24th. When the earlier varieties were being harvested there was no competition other than with those shipped from Baltimore. We believe early cabbage could be made to pay very well, should gardeners concentrate their attention on maturing it a few days earlier than at present. People are always willing to pay more for those home-grown and fresh from the garden.

Select a well-drained and comparatively light soil and give it plenty of manure, at the rate of seventy-five tons to the acre, if possible. The soil should be thoroughly prepared, and to a depth of at least ten inches. Many plow the ground in the fall, which, subjecting it to the freezing and thawing of winter, pulverizes, kills insects, and leaves it in excellent condition to begin with in early Spring. The ground occupied by this crop can be used for a succeeding one, immediately after harvesting. Where the land is very valuable, many also grow radishes, spinach or lettuce between the rows, and by so doing, make these catch crops pay for the manure used. Cuts 3, 4 and 5 show the characteristic forms and comparative sizes of each variety, by a single specimen selected as a representative one. The numbers refer to the same number found in the accompanying table opposite its name.



12

10

19


72

9

2

Cut 4.—First Early Cabbage.

Number.	VARIETY.	Seedsman.	Date of Edible Maturity.	Per Cent.	II aded.	Av. Diam., in.	Av. Weight.	Character of form.
1	Early York.....	Landreth.....	July	10	60	4 3/4	1-4 4-5	See Photograph Cuts 3, 4 and 5.
2	Bloomsdale Bullock Heart.....	Landreth.....		25	72	3 1-5	1-8	
3	Select Very Ey. Jersey Wakefield.....	Landreth.....		28	68	4 1-10	1-8	
4	Louisville Ey. Drumhead.....	Landreth.....		11	44	7 11	1-6 2-5	
5	Landreth's Earliest.....	Landreth.....	June	28	80	5	1-14 2-5	
6	Redlands Ey. Drumhead.....	Landreth.....	July	11	40	5 1/2	1-5	
7	Landreth's Ey. Summer Flathead.....	Landreth.....	June	25	80	5 2-5	1-6 2-5	
8	Early Sugar Loaf.....	Landreth.....		28	68	4	1-6 2-5	
9	N. Jersey Wakefield.....	Burpee.....		25	78	4 1-5	1-3 1-5	
10	Extra Early Express.....	Burpee.....		25	81	4 7-10	1-8	
11	Early Erfurt.....	Burpee.....		28	56	4 2-5	1-2 2-7	
12	Ey. Jersey Wakefield.....	Dunkirk.....		25	96	4 2-5	1-8	
13	Felder Kraut.....	Dunkirk.....	July	11	53	3 1/2	1-1 3-5	
14	Charleston Wakefield.....	Henderson.....	June	24	54	4 2-13	1-4 4-5	
15	Early York.....	Ferry.....	July	8	58	4 1/2	1-4 4-5	
16	Buckbee's New Green.....	Buckbees.....	June	28	68	6	2-0	
18	Landreth's Large York.....	Landreth.....	July	8	68	3 1/2	1-0	
19	Oxheart (French).....	Landreth.....	June	25	88	4 9-10	1-6 2-5	
20	Reynolds.....	Livingston.....		28	92	5	1-6 2-3	
21	Early Jersey Wakefield.....	H. A. March.....		28	96	4	1-6 2-5	
72	Henderson's Ey. Summer.....	Ferry.....		25	96	5 7-10	1-12 4-5	
77	Burpee's Allhead Extra Ey.....	Burpee.....	August	1	80	4 2-5	1-8	



21

20

16

14

11

8

5

3

Cut 5.—First Early Cabbage.

Second Early Cabbage.

Of the *Second Early Cabbage* eighteen varieties were used. All these did fairly well, considering the dry weather. *All Seasons*, N. B. G. & Co., and *Succession*, Henderson, produced heads averaging exactly the same, and were ready for market at the same date. The *All Seasons*, however, produced a slightly greater percentage of matured heads. Following these, although not as large, come *Early Dwarf Flat Dutch*, Ferry, and *Bloomsdale Early Market*, Landreth.


Late Cabbage.

The late varieties of cabbage were not up to the usual standard. Twenty-three varieties were used. The variety averaging the best from all standpoints was the *Early Deep Head*, Livingston. Why this is called *early* we do not know. It makes a solid, oval head. *Christmas King*, Buckbee, is a close competitor and resembles the former very much. Next to these, although not as good, come *Danish Round Winter*, Vick's Sons, *Sure Head*, Burpee, and *Autumn King*, Henderson. The last three were all on a par.

Celery.

This is a vegetable that should be found in every garden. At present nearly all our small towns are furnished by importation. Celery is not difficult to grow; the main thing is to select a comparatively level spot in the garden and to enrich it with well-rotted manure. The plants are generally started from seed in trays or a small bed made for that purpose, and then transplanted into rows in the garden, about six inches apart. In order to make the plants stocky the tops are clipped back when young. It is always desirable to have the soil as moist as possible. When the plants are from ten to twelve inches high, they can be blanched for use; this is accomplished by banking them with loose soil and continuing to do so as the plants grow. The *Golden Self-Blanching* is a variety which, as its name implies, has a tendency to blanch itself, and for the average garden, for home use, although not as large as others, is one of the best.

For the past season the test varieties were very good. Cut 6 shows a single characteristic specimen of each of the varieties. The number in the table below refers to its representation in the cut. The plants are average specimens, and the photograph shows their exact condition as they came from the beds.



1 2 3 4 5 6 7 8 9 10
11 12 13 14 15 16 17 19 20 21 22

Cut 6.—Celery.

Number.	VARIETY.	Seedsman.	No. Stalks.	Height, in.	Weight, lbs.	Av. Weight.	Color.
1	Boston Market.....	Landreth.....	20	24	35.5	1-3 $\frac{3}{4}$	White.
2	Golden Dwarf.....	Landreth.....	20	18	29	1-13 5	White.
3	Half Dwarf.....	Landreth.....	20	24	32.3	1-17-10	White.
4	Large White.....	Landreth.....	20	24	33	1-22-5	White.
5	Sandringham Dwarf White.....	Landreth.....	20	25	32.4	1-22-5	White.
6	Paris Golden Yellow.....	Landreth.....	20	24	34.4	1-11 2 5	White.
7	New White Plume Dwarf.....	Dunkirk.....	20	18	17.5	13 4-5	White.
8	Golden Self-Blanching.....	Dunkirk.....	20	12	11.2	9	White.
9	White Plume.....	Vick Sons.....	20	26	33	1-10	White.
10	New Giant Pascal.....	Vick Sons.....	20	28	32.8	1-10 3-10	White.
11	New Rose.....	Vick Sons.....	20	18	18	15	Pink.
12	New Pink Plume.....	Henderson.....	30	11	9.5	7-1 5	Pink.
13	New Rose.....	U. S. Department.....	20	30	36.5	1-13	Pink.
14	Giant Smooth Prague Celeriac.....	Dreer.....	20	18	12.5	9 4-5	Green.
15	Perle le Grand.....	Livingston's Sons.....	20	24	28	1-6	White.
16	New Perle le Grand.....	Johnson and Stokes.....	20	18	19.5	15 7-10	White.
17	Dwarf Celery.....	Station.....	20	9	7.5	5 4-5	White.
19	Fern Leafed.....	J. L. Childs.....	20	20	28	10 3-10	White.
20	Golden Self-Blanching.....	J. L. Childs.....	20	16	17.2	13 4-5	White.
21	Kalamazoo.....	N. B. G. Co.....	20	18	18.2	14 $\frac{1}{2}$	White.
22	Hartwell Perfection.....	Landreth.....	20	20	34	11-11	White.

Cucumbers.


This vegetable is so commonly grown that little need be said about it other than to name some of the best varieties. The greatest pest we have to contend with in raising the cucumber is the striped beetle. We were able to keep these insects in check by the use of sufficient tobacco dust, which, when dusted upon the young plants, repels them. After the plants get a fair start they will take care of themselves. Twenty-one varieties were grown. For general use the *White Wonder*, Burpee, is one of the best. The plants were very strong as well as productive. The fruit ran very evenly in color, shape and size. The *White Spine*, Landreth, was good for the season, as was the same variety from other seedsmen. *Meaux Green Pickling*, Burpee, new variety; vines very thrifty; fruit very large, July 23rd to September 20th; promising. *Long Green*, Ferry, is one of the largest late varieties. The earliest, although not as good otherwise, was the *Early Cluster*, Landreth.

Egg Plant.

Eight varieties were grown. Many people in West Virginia have never tasted this vegetable, let alone growing it. The Egg Plant is easily grown and a few plants, if in rich soil, will produce enough eggs for a family. They must not be started too early, as they are natives of the tropics, and therefore require rather warm weather. The fruit is fit for use from the time it is the size of a pear until it is fully grown. It is not as good when it indicates an appearance of ripening. The *New York Improved Large Purple* and the *Black Pekin* are two of the best varieties, and are offered by most seedsmen.

Kale.

Five varieties were used in this test, and in cut 7 is represented a characteristic leaf of the following varieties: (1) Scotch Curled, (2) Dwarf Curled Scotch, (3) Dwarf German Greens, (4) Bloomsdale's Extra Doubled Curled, (5) German Dwarf (Purple), (6) Dwarf Curled German. For rank growers, numbers 1, 2 and 5 are preferable; but for dwarf and finely crinkled leaves, numbers, 3, 4 and 6. Culture, similar to that of cabbage. It is used for greens.



Cut 7.—Kale.

Kohl-Rabi.

Three varieties of this vegetable were grown in our trial grounds mainly for the purpose of giving visitors an opportunity to see it. It is not in general use and the sale is very limited. It is sometimes called the turnip-rooted cabbage from resemblance to both the cabbage and turnip. If allowed to get old it becomes stringy and worthless.

Lettuce.

Forty-nine varieties of lettuce have been tested, both in the garden and greenhouse. Of these we prefer the *Grand Rapids*, *Black-Seeded Simpson*, *Hanson* and *Boston Curled* as the best of the open-headed, crinkly varieties. Of the heading varieties, *Frankfort Head* and *Always Ready* were of the best.


The preferable way to grow all kinds of lettuce is to sow the seed thickly and then transplant about ten to twelve inches apart in the row. Beautiful, large heads are thus produced.

Wheeler's Tom Thumb, Burpee, is a variety well named, for it takes but little space in the garden. It is small, forms a very hard, compact head, can be planted close together and matures very early. We also forced it in the greenhouse last season with very good results.

Musk Melon.

Twenty-nine varieties were grown. Among those giving the best satisfaction were *Extra Early White*, Johnson & Stokes, this being small and somewhat flattened on the ends, good quality and quite early; *Banquet*, Barteldes, good size, early, ripening with *Extra Early*, comparatively round, netted, thick yellow meat, small seed cavity, full of seed. *Irondequoit*, Burpee, perfectly smooth, medium to large, white when green, yellowish when ripe, deep yellow flesh, good flavor, September 1st. *Jenny Lind*, Landreth, early, small to medium size, flattened at the poles, good quality. *New Tip Top*, Livingston, good size, white netted, yellowish when ripe, productive; the only objection is that it cracks in the sections when ripe, flesh yellow and sweet, September 10th. *Extra Early Hackensack*, Landreth, medium size, very productive, heavily netted, flesh green, juicy, rich, September 2nd.

Two other varieties possessing merit are the *Jersey Belle* and *Miller Cream*, Johnson & Stokes. These varieties were not in our test, but we have seen them growing and gardeners pronounce them among the best.



Cut 8.—Ordinary and New Culture of Onions.

Onions.

Thirty-five varieties were sown for this test. The season, however, was very dry and no trustworthy results were secured other than with those transplanted. The culture of the onion is ordinarily to sow as early in the Spring as practicable in rich, light soil, in drills about fifteen inches apart, and when the plants are up, thin to three or four inches. Hoe frequently to keep down the weeds. During the past two seasons we found that raising onions with ordinary culture on high land was a failure. The dry season caught them before they were sufficiently grown to withstand it.

The *new method* of onion culture is to sow the seed early in the green house or hot bed, and then to transplant them into the open ground, three or four inches apart, and not too deep. The points claimed for this are that it invariably increases the yield, the onions are larger and more uniform in size, and the extra labor involved in transplanting is offset by the saving of labor in weeding.

Cut 8 represents twelve specimens of the best onions of two varieties, the *Prizetaker*, Barteldes, and *Early Red Globe*, Livingston, treated according to each of the above methods of culture. Number 31 represents twelve of the *Prizetaker* grown according to the new culture; while the small pile to the left represents the best twelve of the same variety under ordinary culture, the seed being sown at the same time that the former were transplanted and coming from the same package. Number 27 represents the *Early Red Globe* under precisely the same conditions as the above.

The following *conclusion*, therefore, can be drawn:

Onions may be grown on high ground in a dry season by the new culture, while under the ordinary culture, it would be a complete loss of seed, time, money and use of land.

Peas.

The following varieties were planted April 21st, in rows five feet apart. They were trained up on a two and one-half foot poultry wire trellis. One hundred vines of each variety were taken as a basis of comparison. The yield, as heretofore given with other vegetables, is on a scale of 0 to 10.

CULTURE—For first early crop sow as early as the ground can be worked, and at intervals of two weeks for succession. Sow in rows two inches deep and from three to four feet apart, according to the variety used and the strength of soil. Extra early varieties sown in August often bring a profitable crop by fall.

Number.	VARIETY.	SEEDMAN.	Per Cent of Germination	Height of Vines. in.	No. Days to Edible Maturity.	Average No. Peas Per Pod.	Yields Scale 0-10, 100 Vines.
1	American Wonder.....	Landreth.....	50	4	59	3	7
2	Landreth's "Extra Early".....	Landreth.....	95	20	58	5	7
3	Alpha.....	Landreth.....	85	23	59	5	7
4	Everbearing.....	Landreth.....	87	23	62	6	7
5	Stratagem.....	Landreth.....	50	24	67	6	9
6	Premium Gem.....	Landreth.....	50	24	62	5	10
7	McLean's Advancer.....	Landreth.....	63	22	62	6	8
8	Pride of the Market.....	Landreth.....	37	25	59	9	10
9	McLean's Little Gem.....	Landreth.....	25	21	59	6	9
10	Telephone.....	Landreth.....	50	29	69	5	8
11	Dw. Blue Imperial.....	Landreth.....	60	21	60	4	8
12	Champion of England.....	Landreth.....	33½	28	70	5	9
14	Days & Summer.....	Landreth.....	100	20	69	5	8
15	Irish Lg. Wht. Marrowfat.....	Landreth.....	100	40	69	5	9
16	The Charmer.....	Vicks & Son.....	75	40	66	7	8
17	Vick's King of the Dwarf.....	Vicks & Son.....	76	14	54	6	8
18	General Grant.....	Buckbee.....	87	35	60	6	9
19	Buckbee's Lightning Express...	Buckbee.....	100	33	57	6	7
21	Deer & Europa.....	Dreer.....	100	16	57	4	6
23	William Hurst.....	Burpee.....	62	9	54	4	7
24	Renown.....	Burpee.....	55	24	69	6	7
25	The Echo.....	Burpee.....	74	23	73	5	8
26	Heroine.....	Henderson.....	76	25	64	8	9
27	Queen.....	Henderson.....	55	24	72	7	7
28	Livingston's First in Market...	Livingston.....	100	25	59	4	7
30	Melting Sugar.....	U. S. Dept.....	80	36	60	7	8
31	Prince of Wales.....	N. B. G. Co.....	85	18	58	5	8
32	Ex. Ey. Summit.....	N. B. G. Co.....	100	24	57	4	7
33	N. B. G. Co's Sapphire.....	N. B. G. Co.....	100	24	57	6	7
35	Sapphire Hero.....	Landreth.....	50	20	64	7	8
36	Ey. Green Marrow.....	Landreth.....	90	16	46	3	6

Peppers.

Nine varieties were used. The cultivation and handling of these plants are precisely similar to that of the Egg Plant, mentioned elsewhere. The only addition to the old Bull Nose type worthy of mention is the *Giant Yellow King*, Livingston. This is, regardless of color, identical with the *Ruby King*. Its being yellow, however, makes it a new addition to our list.

All of the Bull Nose types of peppers were affected by a *Macrosporium* disease, more commonly known as black mould. The fruit began first to decay at the tip, which gradually worked toward the stem end until the whole was worthless. It is thought that perhaps the Bordeaux mixture will keep it in check; the first appearance is somewhat similar to that of the *Macrosporium* of the tomato.

Radishes.

The Radish is especially desired in early Spring. They are easily grown, mature in a short time, and can be used as a catch crop at any place in the garden. They are often planted between rows of beets, cabbage, cauliflower, etc., and are marketed and sold before these crops demand the intervening space. By sowing at intervals of two weeks apart, they may be had fresh for the table or market, throughout the growing season. The best variety for one gardener may not be the best for another, as the choice of color, form, size, etc., depends upon the consumer; the red, however, appears to be the more popular color. The accompanying table shows their comparative merits. Each variety was given a continuous row of eighteen feet. The seed was planted on April 27th.

Number.....	VARIETY.	Seedsman.	Date of Edible Maturity.	Character of Forms.	Color of skin.	Color of Flesh.	Quality.	Average diameter.	Average Weight	Per ct. sound June 8.
1	White tipped Scarlet Turnip...	Landreths.	May 26	Round...	Pink...	White	91.16	.76	40	
2	Ey Scarlet Pruss an Globe.....	Landreths.	22	Round...	Red....	Pink..	91.42	1.33	35	
3	Earliest Scarlet Erfurt.....	Landreths.	23	Turnip...	Red....	White	91.50	1.33	50	
4	Ey Garnet Turnip	Landreths.	23	Round...	D'k red	White	101.42	1.60	30	
5	Summer White	Landreths.	June 10	Conical...	White..	White	72.33	6.40	90	
6	Stutzard White Turnip.....	Landreths.	10	Conical...	White..	White	72.17	4.40	100	
7	Go den Globe	Landreths.	May 26	Conical...	Russett	White	61.33	1.60	95	
8	Ey Deep Scarlet Turnip Rooted	Landreths.	22	Turnip...	D'k red	White	91.50	1.33	40	
9	Ey Red Scarlet Rooted.....	Landreths.	22	Turnip...	D'k red	White	91.33	.80	25	
10	Ey White Turnip	Landreths.	23	Turnip...	White..	White	71.50	1.06	75	
11	New Ey Short Leafed.....	Landreths.	22	Half long	Scarlet	White	71.10	1.12	14	
12	Short Topper Eys't White.....	Landreths.	22	Turnip...	White..	White	91.67	1.60	75	
13	New Crystal Forcing.	Landreths.	28	Conical...	Russett	White	81.16	1.06	100	
14	New Rapid Forcing.....	Buckbee...	22	Round...	Red....	White	71.42	1.12	8	
15	Golden Dresden.....	Burpee	22	Half-R'd.	Russett	White	61.50	1.6	60	
16	French Breakfast.....	Burpee	22	Half long	Red....	White	91.08	.90	15	
17	New Imp. French Breakfast...	Burpee	22	Half long	Red....	White	101.00	.53	20	
18	Earliest Scarlet Button	Burpee	22	Round...	Red....	Pink..	91.50	1.60	60	
19	White Box	N. B. G. Co	30	Turnip.	White..	White	81.67	1.33	70	
20	French Breakfast.....	Landreths...	22	Half long	Red....	White	81.08	.80	20	
21	Burpee's Surprise.....	Burpee	26	Half long	Russett	White	71.42	1.60	90	
22	White Strasberg	N. B. G. Co	26	Long	White..	White	62.17	11.20	95	
24	French Cafe	Landreth...	22	Long	Red....	White	81.17	2.00	30	

Salsify.

Four varieties were used; all of which were of about equal value. This vegetable is coming more and more into use, and every one should give it a trial. It is easily grown, which fact, we feel, is not generally known. The seed is sown in early Spring in drills. The plants need thinning out, should they come up thick. They are not hurt by frost, and can remain in the soil all Winter if desired, and then be dug out in Spring. Some dig them late in the Fall and store them in the cellar, or in a trench outside, to be used at will throughout the season. It is sometimes called the Oyster Plant, from its resemblance in flavor to that of the Oyster.

Summer Squash.

The squash is comparatively a troublesome vegetable to raise on account of its numerous insect enemies. The varieties named in the table were badly infested, which, together with the dry summer, materially lessened the yield. We give the following table, which will indicate the period of ripening, character of form, color of skin, etc. The seed was sown on May 18th. With vegetables that are hard to grow, such as the Summer Squash, this is the time when it pays to take pains with it, on account of the demand and better prices.

Number.	VARIETY.	Seedsman.	Date of Maturity.	Per Cent. of Ma- turity.	No. of Fruits.	Av. Weight, lbs.	Av. Diam., inch.	Character of Form.	Color of Skin.
7	Ex. Ey. Bush.	Landreth.....	July 30	100	17	1.9	8	Knotty, saucer shape.	White.
8	Ey. White Bush.	Landreth.....	July 27	90	16	1.6	7½	Knotty, saucer shape.	White.
10	Long. Green Summer Crook-neck.....	Landreth	Aug. 22	50	15	2.0	4	Long crook- neck.	Green.
11	Landreth's Wht Turban	Landreth	Sept. 20	100	3	1.5	5½	Conical.	White.
12	Golden Summer Crook- neck.	Landreth	July 27	100	25	.9	3½	Long crook- neck.	Yellow
14	Cocozelle.....	Buckbee.	Aug. 22	60	3	4.7	3½	Long.	Green.
15	Silver Custard.	Henderson	Aug. 1	100	10	1.5	7	Round ridged	White.
16	Henderson's Delicata...	Henderson	Aug. 22	100	17	1.2	3½	Long ridged.	Green.
17	Island Prize.....	Johnson & Stokes.	Aug. 16	40	3	1.7	4	Oval.	Yellow.
20	The Der Wing Squash...	Burpee	Aug. 6	100	43	.75	3½	Oval. knotty.	White.

Sweet Corn.

Each variety occupied a row seventy-five feet in length and received ordinary field cultivation. The seed was sown on April 28th. As the table will show, the seed in nearly all cases was deficient in percentage of germination, and the season was so extremely unfavorable for those that did germinate that our gardeners were unable to furnish sufficient sweet corn for Morgantown market. In the column marked "Yield," the scale is 0 to 10; 10 representing the best, 9 good, 8 fair, 7 and below, poor.

Number.	VARIETY.	Seedsman.	Date of Edible Maturity.	Per Cent. of Germination.	Per Cent. of Maturity.	Height of stalk Inches.	Average Length Ears, Inches.	Average Weight Ears, Ounces.	Yield Scale, 0-10.
1	Early Crosby Sugar.	Landreth..	Aug. 13..	53	50	40	6	9.6	5
2	Extra Early Concord Sugar.....	Landreth..	Aug. 13..	16	65	50	6½	8	6
3	Asylum Sugar.	Landreth..	Aug. 2...	35	100	60	6	6.4	7
4	Landreth's Sugar.....	Landreth..	Aug. 13..	45	70	60	7	11.2	7
5	Evergreen Sweet Corn	Landreth..	Aug. 2...	35	71	60	7	8	6
6	Late Mammoth Sugar.....	Landreth..	Aug. 2...	36	100	62	8	9.6	7
7	Early Mammoth Sugar.....	Landreth..	Aug. 2...	36	100	62	8	8	8
14	Extra Early Adams	Landreth..	July 14..	55	86	65	6½	8	8
15	Early Landreth Market.	Landreth..	July 30..	69	95	72	8	8	8
16	Extra Early Minnesota Sugar	Landreth..	July 15..	35	35	60	6½	6.4	5
17	Extra Early Naragansett Sugar.....	Landreth..	July 16..	41	39	46	6	4.8	6
18	Buckbee's "Best of All".....	Buckbee..	July 25..	40	43	56	6	6.4	6
19	Port and	N. B. G. Co	July 25..	49	58	58	6	6.4	7
20	Zig Zag Evergreen.....	N. B. G. Co	Aug. 17--	41	72	60	6½	9.6	8
21	Early Ford Hook.....	Burpee ..	July 14..	36	79	46	6½	4.8	7
22	Livingston's New Silver Cane.	Livingston	Aug. 13..	35	95	52	8	12.8	7
23	Zig Zag Evergreen.....	U. S. Dept.	Aug. 6...	66	100	61	7	8	9
26	Red Cob Corn.....	Station ..	Aug. 22..	98	37	68	6	11.2	10
27	Dungan's Sweet Corn.....	Barteldes..	Aug. 30..	100	48	68	7	11.2	8
28	Country Gent.....	Barteldes..	Aug. 11..	73	50	74	6	6.4	7

Tomatoes.

The seed was sown in the forcing house, March 12th, in ordinary trays, and transplanted and placed in cold frames before going into the trial grounds on May 8th. Ten plants were taken as the test of each variety. The rows were five feet apart, and the plants three and one-half feet apart in the row. Each plant was tied to a stake to keep the vines in place. A small black flea beetle was very troublesome while the plants were young. The following table will point out the comparative results; the headings of the columns will explain themselves.

No.	VARIETY.	SEEDSMAN.	Date of First Ripeness	Yield of Ripe Fruit to Aug. 15th.		Yield from Aug. 15th to Sept. 1.		Yield from Sept. 1 to Sept. 26.		Total Weight Ripe Fruits.		Average Weight Ripe Fruits.		Weight Green Part left on Vines.		Total Weight Rotten Fruits.		Total Weight.		REMARKS.
				Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Oz.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	
1	Extra Early Jersey.....	Landreth.....	July	30	19.9	25.1	23.3	23.3	61.3	4.80	10.5	3.1	71.8	10.5	3.1	81.8	3.1	71.8	3.1	Early but rough.
2	Early Richmond.....	Landreth.....	July	27	37.8	19.2	28.7	28.7	85.7	3.20	21.3	3.6	108.3	21.3	3.6	108.3	3.6	108.3	3.6	Good, Early.
3	Beauty Second Early.....	Landreth.....	July	26	6.1	26.9	26.6	26.6	59.6	3.20	21.3	3.6	81.8	21.3	3.6	81.8	3.6	81.8	3.6	Good, Early.
4	Bermuda Extra Early.....	Landreth.....	July	26	35.8	22.3	31.5	31.5	89.6	3.20	21.3	3.6	108.3	21.3	3.6	108.3	3.6	108.3	3.6	Good, Early.
5	Hathaway's Excelsior.....	Landreth.....	July	30	8.1	42.1	35.2	35.2	85.4	3.20	21.3	3.6	108.3	21.3	3.6	108.3	3.6	108.3	3.6	Good, Early.
6	Trophy.....	Landreth.....	July	27	19.6	20.0	20.0	20.0	76.5	3.20	21.3	3.6	108.3	21.3	3.6	108.3	3.6	108.3	3.6	Good, Early.
7	Go den Trophy.....	Landreth.....	July	30	4.1	20.4	20.4	20.4	76.5	3.20	21.3	3.6	108.3	21.3	3.6	108.3	3.6	108.3	3.6	Good, Early.
8	Cardinal.....	Landreth.....	July	26	18.5	25.5	26.3	26.3	70.3	3.20	21.3	3.6	108.3	21.3	3.6	108.3	3.6	108.3	3.6	Good, Early.
9	Perce Improved.....	Landreth.....	July	31	5.5	14.7	29.2	29.2	44.4	4.48	15.5	7.0	59.9	15.5	7.0	59.9	7.0	59.9	7.0	Good, Early.
10	Livingston's Favorite.....	Landreth.....	July	24.1	28.9	20.6	20.6	20.6	73.6	2.88	18.4	13.2	91.1	18.4	13.2	91.1	13.2	91.1	13.2	Good, Early.
11	Dwarf Champion.....	Livingston's Sons.....	August	10	1.4	12.1	16.4	16.4	29.8	2.56	7.0	3.4	36.8	7.0	3.4	36.8	3.4	36.8	3.4	Good, Early.
12	Livingston's Beauty.....	Livingston's Sons.....	August	30	16.6	31.9	27.4	27.4	75.9	2.56	7.0	3.4	36.8	7.0	3.4	36.8	3.4	36.8	3.4	Good, Early.
13	Bu'keye State.....	Livingston's Sons.....	July	22	5.5	3.1	45.1	45.1	48.2	4.80	16.7	3.3	70.7	16.7	3.3	70.7	3.3	70.7	3.3	Good, Early.
14	Livingston's Gold Ball.....	Livingston's Sons.....	July	27	2.5	18.5	30.1	30.1	54.1	3.84	9.0	1.0	56.5	9.0	1.0	56.5	1.0	56.5	1.0	Good, Early.
15	Dwarf Aristocrat.....	Livingston's Sons.....	August	1	2.5	13.2	11.4	11.4	47.5	2.72	11.0	9.0	42.8	11.0	9.0	42.8	9.0	42.8	9.0	Good, Early.
16	Paragon.....	Dunkirk.....	August	30	4.3	20.6	22.8	22.8	47.7	2.88	12.0	7.8	59.7	12.0	7.8	59.7	7.8	59.7	7.8	Good, Early.
17	Large Red.....	Dunkirk.....	July	30	7.7	18.6	19.8	19.8	46.1	2.40	12.5	8.5	58.6	12.5	8.5	58.6	8.5	58.6	8.5	Good, Early.
18	Early York.....	Dunkirk.....	July	27	8.9	22.7	23.4	23.4	55.0	3.58	24.0	16.1	79.0	24.0	16.1	79.0	16.1	79.0	16.1	Good, Early.
19	General Grant.....	Dunkirk Seed Co.....	July	27	1.4	13.0	13.7	13.7	28.1	2.08	8.3	8.3	36.4	8.3	8.3	36.4	8.3	36.4	8.3	Good, Early.
20	Canada Victor.....	Ferry & Co.....	July	27	16.3	19.8	26.3	26.3	58.3	3.72	30.0	12.2	78.3	30.0	12.2	78.3	12.2	78.3	12.2	Good, Early.
21	Paragon.....	Ferry & Co.....	July	30	5.3	17.1	22.6	22.6	53.7	1.76	34.9	3.0	42.9	34.9	3.0	42.9	3.0	42.9	3.0	Good, Early.
22	Ex. Early Smooth Red.....	Ferry & Co.....	July	27	9.1	21.5	22.6	22.6	53.7	2.24	13.5	5.6	67.2	13.5	5.6	67.2	5.6	67.2	5.6	Good, Early.
23	Livingston's Perfection.....	Ferry & Co.....	July	27	14.6	27.5	23.2	23.2	65.3	2.40	18.0	21.5	83.3	18.0	21.5	83.3	21.5	83.3	21.5	Good, Early.
24	The Matchless.....	Albee Rupee & Co.....	July	27	4.0	13.3	17.3	17.3	34.6	2.24	6.3	5.1	40.9	6.3	5.1	40.9	5.1	40.9	5.1	Good, Early.
25	Fordhook First.....	Albee Rupee & Co.....	July	30	4.0	13.3	17.3	17.3	34.6	2.24	6.3	5.1	40.9	6.3	5.1	40.9	5.1	40.9	5.1	Good, Early.
26	Royal Red.....	Bareldes.....	July	30	23.5	24.4	23.7	23.7	71.6	2.56	22.0	13.4	93.6	22.0	13.4	93.6	13.4	93.6	13.4	Good, Early.
27	Buckley's New Tree.....	Buckbee.....	July	33	17.1	13.5	24.2	24.2	54.8	2.24	13.0	8.0	67.8	13.0	8.0	67.8	8.0	67.8	8.0	Good, Early.
28	Crimson Cushion.....	Henderson.....	July	30	18.7	17.1	13.5	13.5	54.8	2.24	13.0	8.0	67.8	13.0	8.0	67.8	8.0	67.8	8.0	Good, Early.
29	Henderson's Ponderosa.....	Henderson.....	August	8	2.1	53.8	42.2	42.2	97.6	7.08	22.0	5.9	89.4	22.0	5.9	89.4	5.9	89.4	5.9	Good, Early.
30	Stanton Free.....	J. L. Childs.....	July	27	15.3	26.6	19.6	19.6	61.5	4.40	31.0	7.4	92.5	31.0	7.4	92.5	7.4	92.5	7.4	Good, Early.
31	Climbing.....	J. L. Childs.....	July	27	11.4	27.2	24.1	24.1	62.7	2.88	22.0	8.8	113.7	22.0	8.8	113.7	8.8	113.7	8.8	Good, Early.
32	Brunton's Best.....	Johnson & Stokes.....	July	27	29.6	38.5	22.6	22.6	90.7	3.90	23.0	9.4	143.7	23.0	9.4	143.7	9.4	143.7	9.4	Good, Early.
33	New Liberty Bell.....	Johnson & Stokes.....	July	27	3.8	34.0	42.6	42.6	80.4	6.56	40.0	16.9	128.7	40.0	16.9	128.7	16.9	128.7	16.9	Good, Early.
34	Ring Leader.....	Dreer.....	August	13	27	42.2	35.3	35.3	108.7	2.40	19.0	10.2	96.2	19.0	10.2	96.2	10.2	96.2	10.2	Good, Early.
35	Ponderosa.....	U. S. Dept.....	July	27	24.7	27.4	25.1	25.1	77.2	2.08	19.0	10.2	96.2	19.0	10.2	96.2	10.2	96.2	10.2	Good, Early.
36	Acme.....	Ferry & Co.....	July	30	24.7	27.4	25.1	25.1	77.2	2.08	19.0	10.2	96.2	19.0	10.2	96.2	10.2	96.2	10.2	Good, Early.

Smooth and Uniform.

Smooth, Round and Uniform.

Smooth and Uniform.

Smooth, Round and Uniform.

Smooth and Uniform.

Smooth, Round and Uniform.

Number.	VARIETY.	Seedsman.	Date of First Ripe Fruit.	Yield of Ripe Fruit to Aug.		Yield from September 1 to September 26.		Total Weight of Ripe Fruits.		Average Weight of Ripe Fruits.		Weight of Green Fruit Left on Vines.		Total Weight of Rotten Fruit.		Total Weight of Fruit Grown.		Remarks.
				Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Oz.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	
45	Golden Fig	J. L. Childs	July	30	24.3	43.1	25.6	93.0	0.32	13.0	0	11.4	0	106.0				Inferior quality.
47	Ten Ten Tomato	Landreth		30	16.9	25.7	22.7	65.3	2.40	21.0	11.4	25.0	11.4	86.3				
48	Chemine Apple	Landreth		30	18.0	42.7	26.0	86.7	3.20	38.0	16.9	38.0	16.9	111.7				Large, uniform, smooth.
49	Buckeye	Landreth		27	8.7	24.6	53.0	85.3	3.84	38.0	1.8	38.0	1.8	123.3				
50	Baltimore Prize Taker	Landreth		27	33.8	49.3	33.9	117.0	2.88	30.0	19.8	30.0	19.8	147.0				
51	Atlantic Prize	Landreth		27	21.2	15.0	27.1	63.3	2.54	20.0	1.1	20.0	1.1	83.3				
52	Money Maker	Landreth		30	26.8	26.2	36.7	88.7	4.48	35.3	8.5	35.3	8.5	115.3				
53	May's Favorite	R. D. Hawley & Co		23	26.9	30.8	24.5	82.2	3.20	27.0	18.8	27.0	18.8	109.2				
54	Stone	Landreth	August	3	1.5	20.3	20.2	42.0	4.00	16.0	16.2	16.0	16.2	58.0				Rough and inferior.
55	World's Fair	Read	July	30	19.9	30.9	31.1	81.9	1.92	21.0	1.8	21.0	1.8	102.2				
56	Cross No. 1	B. W. Alwood		27	24.0	34.6	27.2	85.8	4.16	24.0	18.5	24.0	18.5	103.8				
57	Toss No. 2	B. W. Alwood		30	16.1	52.7	34.8	103.1	3.20	32.0	17.4	32.0	17.4	135.1				
58	Cross No. 3	B. W. Alwood		30	18.2	36.6	27.5	82.3	3.20	26.0	19.9	26.0	19.9	108.3				
59	Cross No. 4	B. W. Alwood		27	27.2	49.2	23.6	106.0	3.04	38.0	29.0	38.0	29.0	144.0				
60	Lemon Blush	Station	August	1	26.0	51.7	38.0	115.7	2.72	31.0	11.5	31.0	11.5	146.7				
61	Seedling No. 1	Dr. Casselberry		6	3.5	34.9	64.1	102.5	6.88	30.0	1.1	30.0	1.1	132.5				Largest 1.2 lb.
62	Terra Corta	Station	July	18	48.2	34.7	24.9	107.8	2.56	59.5	10.2	59.5	10.2	129.3				Soft and seedy.
63	Seedling No. 2	Station	August	3	11.7	54.6	23.9	95.2	8.96	59.5	3.1	59.5	3.1	151.7				
64	Seedling No. 3	Station		6	11.0	46.7	23.1	80.8	8.48	56.0	3.1	56.0	3.1	136.8				
65	Seedling No. 4	Station		13	5.0	25.0	40.0	70.0	9.60	54.5	0	54.5	0	131.5				

Water Melon.

Fifteen varieties were used. The old standards are too well known to require special mention. Two new varieties not so commonly grown, worthy of mention are: *Fordhook Early*, Burpee, very early, round in form, a medium green, striped; flesh, bright red with white seed, good size and quality. *Kentucky Wonder*, N. B. G. & Co., large, oblong in form, skin medium green, flesh red, seeds brown, good quality.

Miscellaneous.

Numerous other kinds of vegetables were grown, as *Parsnips*, *Brussels—Sprouts*, *Borocoli*, *Turnips*, *Carrots*, *Collards*, *Cress*, *Endive*, *Leek*, *Mustard*, *Okra*, *Parsley*, *Winter Radishes*, *Ruta-Bagas*, *Spinach*, *Rhubarb*, etc. Among these there are comparatively few new introductions; therefore, we do not deem it necessary to give any detailed account of each.

Donations to the Horticultural Department, 1894.

Names.	Pkt's Flow. Seed	Pkt's Veg. Seed
J. A. Everitt.....	5
H. W. Buckbee.....	31
W. Atlee Burpee.....	9	19
Prof. W. B. Alwood.....	4
Jas. Vick & Sons.....	5	4
Northup, Braslon, Goodwin & Co.	5	8
H. A. March.....	4
L. H. Reid.....	1
Prof. L. H. Bailey.....	4
Jas. M. Thornburn & Co.....	3
A. W. Livingston's Sons.....	3
H. A. Dreer.....	5
Peter Henderson & Co.....	1	12
U. S. Department Agriculture...	22

Remarks.

1. The Table of Contents, page 137, refers you to the various kinds of vegetables treated.

2. If any one desires information in regard to any variety not mentioned in this Bulletin, we will be pleased to aid them, if possible.

3. Our purpose at present is to test all the new introductions and to report upon their comparative merits, that we may aid in the selection of seeds.

4. The conclusions in tests are based upon both our own experiments and those of other noted and trusted gardeners.

F. WM. RANE.



